


Navigating the December Night Sky

For observers in the middle northern latitudes, this chart is suitable for late November at 9 p.m. or early December at 8 p.m.

The stars plotted represent those which can be seen from areas suffering from moderate light pollution. In larger cities, less than 100 stars are visible, while from dark, rural areas well over ten times that amount are found.



Navigating the late fall night sky: Simply start with what you know or with what you can easily find.

- 1 Face south. Almost overhead is the "Great Square" with four stars about the same brightness as those of the Big Dipper. Extend an imaginary line southward following the Square's two westernmost stars. The line strikes Fomalhaut, the brightest star in the southwest. A line extending southward from the two easternmost stars, passes Deneb Kaitos, the second brightest star in the south.
- 2 Draw another line, this time westward following the southern edge of the Square. It strikes Altair, part of the "Summer Triangle."
- 3 Locate Vega and Deneb, the other two stars of the "Summer Triangle." Vega is its brightest member while Deneb sits in the middle of the Milky Way.
- 4 Jump along the Milky Way from Deneb to Cepheus, which resembles the outline of a house. Continue jumping to the "W" of Cassiopeia, to Perseus, and finally to Auriga with its bright star Capella.

Binocular Highlights

- A and B:** Examine the stars of the Pleiades and Hyades, two naked eye star clusters.
- C:** The three westernmost stars of Cassiopeia's "W" point south to M31, the Andromeda Galaxy, a "fuzzy" oval.
- D:** Sweep along the Milky Way from Altair, past Deneb, through Cepheus, Cassiopeia and Perseus, then to Auriga for many intriguing star clusters and nebulous areas.

