


# Navigating the June Night Sky

For observers in the middle northern latitudes, this chart is suitable for early June at 11:30 p.m. or late June near 10:30 p.m.

The stars plotted represent those which can be seen from areas suffering from moderate light pollution. In larger cities, less than 100 stars are visible, while from dark, rural areas well over ten times that amount are found.



**Navigating the mid June night sky: Simply start with what you know or with what you can easily find.**

- 1 Extend a line north from the two stars at the tip of the Big Dipper's bowl. It passes by Polaris, the North Star.
- 2 Draw another line in the opposite direction. It strikes the constellation Leo high in the west.
- 3 Follow the arc of the Dipper's handle. It first intersects Arcturus, the brightest star in the June evening sky, then Spica.
- 4 Arcturus, Spica, and Denebola form the Spring Triangle, a large equilateral triangle.
- 5 To the northeast of Arcturus shines another star of the same brightness, Vega. Draw a line from Arcturus to Vega. It first meets "The Northern Crown," then the "Keystone of Hercules." A dark sky is needed to see these two dim stellar configurations.
- 6 High in the east are the three bright stars of the Summer Triangle: Vega, Altair, and Deneb.

## Binocular Highlights

- A: Between Denebola and the tip of the Big Dipper's handle, lie the stars of the Coma Berenices Star Cluster.
- B: Between the bright stars of Antares and Altair, hides an area containing many star clusters and nebulae.
- C: 40% of the way between Altair and Vega, twinkles the "Coathanger," a group of stars outlining a coathanger.
- D: Sweep along the Milky Way for an astounding number of faint glows and dark bays.

